



**CORSO**  
**AGGIORNAMENTO PER MICOLOGO**  
**ORE 12**

- Cortinarius subgen. Phlegmacium (specie commestibili a carne bianca e gambo privo di bulbo basale marginato (C. praestans, C. largus, C. cumatilis, C. balteatocumatilis, C. claricolor, C. varius, C. caligatus, ecc.)
- Lactarius sez. Dapetes (L. deliciosus, L. salmonicolor, L. deterrimus, L. sanguifluus, L. semisanguifluus, ecc.)
- Boletus sez. Boletus (Boletus s.str.) (B. edulis, B. pinophilus, B. reticulatus, B. aereus)
- Armillaria (A. mellea, A. caepistipes, A. gallica, A. ostoyae) + Desarmillaria (D. tabescens)
- Cantharellus (C. cibarius, C. friesii, C. pallens, C. ferruginascens, C. alborufescens, C. amethysteus) + Craterellus (C. lutescens, C. tubaeforme, C. cornucopiae, C. cinereus, C. melanoxeros, C. ianthinoxanthus) + Pseudocraterellus (P. sinuosus)
- Russula (Plorantinae) (R. delica, R. chloroides) (R. cyanoxantha, R. vesca, R. heterophylla, R. mustelina, R. aurea, ecc.)
- Hygrophorus (H. marzuolus, H. penarioides, H. russula)
- Altre specie single distribuite in vari generi